


# CARBOHYDRATE DERIVED LIGANDS FOR CATALYTIC STEREOSELECTIVE CARBENOID CYCLOPROPANATION REACTIONS

Felix Preusch and Thomas Ziegler

Institute of Organic Chemistry, University of Tuebingen, Auf der Morgenstelle 18,  
72076 Tuebingen, Germany, thomas.ziegler@uni-tuebingen.de

Metal-catalyzed stereo selective cyclopropanation plays an important role in organic synthesis.[1] Stereoselective introduction of a cyclopropane unit to an intermediate together with a vast number of ring-opening possibilities lead to an important tool in precursor synthesis.[2,3]

It has already been shown that carbohydrate derived bis(oxazoline) ligands can catalyze copper carbenoid cyclopropanations in good enantiomeric excess.[4] Metal-catalyzed cyclopropanation has its origin in a copper catalyzed reaction. Nevertheless, a modern approach to enantioselective cyclopropanation employs binuclear rhodium(II)tetracarboxylate complexes.[5,6]



In this work, we present the synthesis of new carbohydrate derived rhodium carboxylate complexes as well as their application in enantioselective carbenoid cyclopropanation catalysis.

[1] H. Reissig, R. Zimmer, *Chem. Rev.*, **2003**, *103*, 1151–1196.

[2] L. A. Paquette, *Chem. Rev.*, **1986**, *86*, 733–750.

[3] A. de Meijere, L. Wessjohann, *Synlett*, **1990**, 20–32.

[4] M. Irmak, A. Groschner, M. M. K. Boysen, *Chem. Commun.*, **2007**, *0*, 177–179

[5] H. Nozaki, S. Moriuti, H. Takaya, R. Noyori, *Tetrahedron Lett.*, **1966**, *7*, 5239–5244.

[6] A. J. Hubert, A. F. Noels, A. J. Anciaux, P- Teyssié, *Synthesis*, **1976**, *09*, 600–602.