

## A CHIMERA MODEL OF Rel<sub>Seq</sub> PRE-CATALYTIC STATE

Monica Civera and Sara Sattin

Università degli Studi di Milano, Department of Chemistry, Via Golgi, 19, 20133,  
Milan, Italy

Bacteria use nucleotides as second messengers to regulate diverse cellular processes in response to environmental stimuli. For instance, intracellular accumulation of guanosine tetra- or pentaphosphate ((p)ppGpp) leads to a temporary antibiotic-tolerant phenotype called persister, playing a starring role in chronic and recurrent infections [1]. (p)ppGpp is synthesized and/or hydrolyzed by proteins belonging to the RelA/SpoT-Homologue (RSH) superfamily. The multi-domain ‘long’ RSH enzymes, such as Rel<sub>Seq</sub> [2] (*S. equisimilis*), have both a synthetase (SYNTH) and a hydrolase (HD) domain. Along with long RSHs, shorter and specialized RSHs that carry either the SYNTH or the HD domain have been identified. Recently, the X-ray crystal structure of the pre-catalytic state of the *S. aureus* single synthetase RelP was reported [3]. The structural homology between the SYNTH domains of these two proteins led us to consider the pre-catalytic binding site of RelP as a reliable template for building the pre-catalytic state of Rel<sub>Seq</sub>.

Here we report a composite homology model of the bifunctional enzyme generated using both the X-ray structure of Rel<sub>Seq</sub> (to model the HD domain) and RelP (to model the SYNTH pre-catalytic state) as templates. Molecular dynamics simulations were then performed to assess the stability of the model.

---

[1] K. Gerdes and E. Maisonneuve, *Cell*, **2014**, 2, 539-548

[2] T. Hogg *et al.*, *Cell*, **2014**, 117, 57–68

[3] M. C. Manav *et al.*, *J. Biol. Chem.*, **2018**, 293(9), 3254-3264