


# CONCISE SYNTHESIS OF ALL-*CIS*-3,4,5-SUBSTITUTED PIPERIDINES THROUGH CASCADE RADICAL CYCLISATION INITIATED BY VISIBLE LIGHT-PROMOTED HYDROSILYLATION OF ALKENES

Zhu-Jun Yao, Wei-Chen Cui, Shaozhong Wang, and Jing Zhu

State Key Laboratory of Coordination Chemistry, Jiangsu Key Laboratory of Advanced Organic Materials, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing, Jiangsu 210023, China  
yaoz@nju.edu.cn

The piperidine structural motif is widely present in numerous natural alkaloids and marketed pharmaceuticals,<sup>1</sup> and numerous methodologies have been reported to synthesize various types of piperidines. However, synthesis of all-*cis*-multisubstituted piperidines keeps a challenge with current protocols yet. In recent years, several methods on visible light-initiated hydrosilylation of alkenes and alkynes were developed by our group.<sup>2-3</sup> Herein, we wish to report a concise one-step synthesis of all-*cis*-3,4,5-substituted piperidines through cascade radical cyclisation initiated by the visible light-promoted hydrosilylation of alkenes.<sup>4</sup>



[1] R. Vardanyan, Piperidine-Based Drug Discovery, Elsevier Inc., **2018**.

[2] J. Zhu, W.-C. Cui, S. Wang, Z.-J. Yao, *Organic Letters* **2018**, *20*, 3174-3178.

[3] J. Zhu, W.-C. Cui, S. Wang, Z.-J. Yao, *J. Org. Chem.* **2018**, *83*, 14600–14609.

[4] W.-C. Cui, J. Zhu, S. Wang, Z.-J. Yao, unpublished work, **2019**.