

# SYNTHESIS AND CHARACTERIZATION OF SELENOPHENE-BASED ORGANIC SEMICONDUCTORS

Brigitte Holzer<sup>a</sup>, Berthold Stöger<sup>b</sup>, Daniel Lumpi<sup>a</sup>, Ernst Horkel<sup>a</sup>,  
and Johannes Fröhlich<sup>a</sup>

<sup>a</sup>Institute of Applied Synthetic Chemistry, TU Wien (Vienna University of Technology), 1060 Vienna, Austria

<sup>b</sup>X-Ray Centre, TU Wien (Vienna University of Technology), 1060 Vienna, Austria

Thiophene-based small molecules proved to be benchmark p-type organic semiconductors owing to their unique electronic properties associated with their  $\pi$ -electron topology [1]. Recently, several promising organic semiconductors based on benzothieno[3,2-*b*]thiophene **1a** [2] and benzothieno[2,3-*b*]thiophene subunits **2a** [3, 4] (Figure 1) yielding high charge carrier mobilities have been investigated. The introduction of electron donating, and highly polarizable selenium may even enhance charge carrier mobilities. However, the development of reliable synthetic pathways toward selenium-based materials is still matter of ongoing research owing to a lack of available starting materials.

Based on the structural moieties **1a** and **2a**, the topic of this contribution focusses on the integration of selenium in these fused  $\pi$ -conjugated compounds leading to semiconductors **1b-1d** and **2b-2d** (Figure 1).




Figure 1: Mixed thiophene/selenophene fused target compounds.

Reliable protocols towards regioisomeric selenophene-based fused acenes will be presented starting from commercially available building blocks like benzo[*b*]thiophene and benzo[*b*]selenophene. Furthermore, the photo-physical and electro-chemical properties of these target materials will be probed by cyclic voltammetry and UV-Vis spectroscopy.

[1] Q. Meng, et al., J. Mater. Chem. 21, 11708 (2011).

[2] H. Chen, et al., J. Phys. Chem. C 115, 23984 (2011).

[3] T. Mathis, et al., J. Appl. Phys. 115, 043707 (2014).

[4] Y. Liu et al., J. Mater. Chem. C 2, 8804 (2014).