


# $\gamma$ -SUBSTITUENTS AS TOOLS TO INFLUENCE THE REACTIVITY AND STEREOSELECTIVITY OF PROLINE BASED CATALYSTS

Martin Schnurr and Helma Wennemers

Laboratory of Organic Chemistry, ETH Zürich, Vladimir-Prelog-Weg 3, Zürich, 8093, Switzerland

Tripeptides of the H-Pro-Pro-Xaa type are highly reactive and stereoselective catalysts for asymmetric aldol reactions and conjugate addition reactions of carbonyl compounds to nitroolefins, dicyanoolefins and unprotected maleimide [1]. For example, as little as 0.1 mol% H-DPro-Pro-Glu-NH<sub>2</sub> suffices to catalyze conjugate addition reactions of aldehydes to nitroolefins in high yields and excellent stereoselectivities [2]. Mechanistic studies showed, that the catalytic cycle proceeds *via* an enamine intermediate, which takes part in the rate- and stereodetermining C–C-bond formation step [3]. We envisioned that the enamine reactivity and stereoselectivity can be influenced by changing the conformation of the *N*-terminal proline. We introduced substituents at the  $\gamma$ -position of the proline ring, which are known to favor different ring puckers [4]. We used the conjugate addition reaction of butanal to nitrostyrene as testing ground to analyze the performance of the  $\gamma$ -substituted catalysts by *in situ* IR- and NMR-spectroscopic experiments. We complemented our findings by studies on enamine model compounds, which were derived from  $\gamma$ -substituted proline methyl esters. Our investigations provided insight into the effect of different ring puckers on the reactivity and stereoselectivity of proline derived catalysts.



[1] a) P. Krattiger, R. Kovasy, J. D. Revell, S. Ivan, H. Wennemers, *Org. Lett.* **2005**, *7*, 1101–1103. b) M. Wiesner, J. D. Revell, H. Wennemers, *Angew. Chem. Int. Ed.* **2008**, *120*, 1897–1900. c) J. Duschmalé, H. Wennemers, *Chem. Eur. J.* **2012**, *18*, 1111–1120. d) R. Kastl, H. Wennemers, *Angew. Chem. Int. Ed.* **2013**, *52*, 7228–7232. e) T. Schnitzer, H. Wennemers, *Synlett* **2017**, *28*, 1282–1286. f) C. E. Grünenfelder, J. K. Kisunzu, H. Wennemers, *Angew. Chem. Int. Ed.* **2016**, *55*, 8571–8574.

[2] a) M. Wiesner, G. Upert, G. Angelici, H. Wennemers, *J. Am. Chem. Soc.* **2010**, *132*, 6–7. b) T. Schnitzer, H. Wennemers, *J. Am. Chem. Soc.* **2017**, *139*, 15356–15362.

[3] F. Bächle, J. Duschmalé, C. Ebner, A. Pfaltz, H. Wennemers, *Angew. Chem. Int. Ed.* **2013**, *52*, 12619–12623.

[4] a) L. E. Zimmer, C. Sparr, R. Gilmour, *Angew. Chem. Int. Ed.* **2011**, *50*, 11860–11871. b) L. S. Sonntag, S. Schweizer, C. Ochsenfeld, H. Wennemers, *J. Am. Chem. Soc.* **2006**, *128*, 14697–14703.