


NOVEL APPROACH TO HETEROCYCLES BY TRANSITION METAL CATALYZED [2+2+2] CYCLOADDITIONS

Tim Gläsel^{a, b}, Marko Hapke^{a, b}

^aInstitute for Catalysis (INCA), Johannes Kepler University, 4040 Linz, Austria

^bLeibniz Institute for Catalysis e. V. (LIKAT), 18059 Rostock, Germany

Cyclotrimerization reactions have matured in the recent decades to a valuable tool for the construction of carbo- and heterocarbocyclic ring system.[1] Transition metal complexes as suitable catalysts can now be found throughout the whole range of these metals in the periodic table, each possessing some individual properties and advantages.[2] We have specialized in cobalt-catalyzed cyclotrimerization reactions in synthetic heterocyclic chemistry, enabling the construction of complex ring structures from smaller synthons.[3] We are going to present novel heterocycles assembled by the transition metal-catalyzed [2+2+2] cycloaddition employing different metals and thereby utilizing unusual reaction conditions as well as providing the opportunity for facile further modifications of the obtained multifunctional heterocycles.



[1] K. Tanaka (Ed.) (2013). *Transition-Metal-Mediated Aromatic Ring Construction*. Hoboken, USA: John Wiley & Sons.

[2] M. Hapke, N. Weding, K. Kral in *Applied Homogeneous Catalysis with Organometallic Compounds: A Comprehensive Handbook in Four Volumes, Third Edition* edited by B. Cornils, W. A. Herrmann, M. Beller, R. Paciello, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2018, 1, 375-392.

[3] P. Jungk, F. Fischer, M. Hapke, *ACS Catalysis*, 2016, 6, 3025-3029; F. Fischer, M. Hapke, *Eur. J. Org. Chem.* 2018, 3193-3201.